



BRIGHT FUTURES HANDOUT ► PARENT

12 MONTH VISIT

Here are some suggestions from Bright Futures experts that may be of value to your family.

✓ HOW YOUR FAMILY IS DOING

- If you are worried about your living or food situation, reach out for help. Community agencies and programs such as WIC and SNAP can provide information and assistance.
- Don't smoke or use e-cigarettes. Keep your home and car smoke-free. Tobacco-free spaces keep children healthy.
- Don't use alcohol or drugs.
- Make sure everyone who cares for your child offers healthy foods, avoids sweets, provides time for active play, and uses the same rules for discipline that you do.
- Make sure the places your child stays are safe.
- Think about joining a toddler playgroup or taking a parenting class.
- Take time for yourself and your partner.
- Keep in contact with family and friends.

✓ ESTABLISHING ROUTINES

- Praise your child when he does what you ask him to do.
- Use short and simple rules for your child.
- Try not to hit, spank, or yell at your child.
- Use short time-outs when your child isn't following directions.
- Distract your child with something he likes when he starts to get upset.
- Play with and read to your child often.
- Your child should have at least one nap a day.
- Make the hour before bedtime loving and calm, with reading, singing, and a favorite toy.
- Avoid letting your child watch TV or play on a tablet or smartphone.
- Consider making a family media plan. It helps you make rules for media use and balance screen time with other activities, including exercise.

✓ FEEDING YOUR CHILD

- Offer healthy foods for meals and snacks. Give 3 meals and 2 to 3 snacks spaced evenly over the day.
- Avoid small, hard foods that can cause choking—popcorn, hot dogs, grapes, nuts, and hard, raw vegetables.
- Have your child eat with the rest of the family during mealtime.
- Encourage your child to feed herself.
- Use a small plate and cup for eating and drinking.
- Be patient with your child as she learns to eat without help.
- Let your child decide what and how much to eat. End her meal when she stops eating.
- Make sure caregivers follow the same ideas and routines for meals that you do.

✓ FINDING A DENTIST

- Take your child for a first dental visit as soon as her first tooth erupts or by 12 months of age.
- Brush your child's teeth twice a day with a soft toothbrush. Use a small smear of fluoride toothpaste (no more than a grain of rice).
- If you are still using a bottle, offer only water.

Helpful Resources: Smoking Quit Line: 800-784-8669 | Family Media Use Plan: www.healthychildren.org/MediaUsePlan
Poison Help Line: 800-222-1222 | Information About Car Safety Seats: www.safercar.gov/parents | Toll-free Auto Safety Hotline: 888-327-4236

12 MONTH VISIT—PARENT



SAFETY

- Make sure your child's car safety seat is rear facing until he reaches the highest weight or height allowed by the car safety seat's manufacturer. In most cases, this will be well past the second birthday.
- Never put your child in the front seat of a vehicle that has a passenger airbag. The back seat is safest.
- Place gates at the top and bottom of stairs. Install operable window guards on windows at the second story and higher. Operable means that, in an emergency, an adult can open the window.
- Keep furniture away from windows.
- Make sure TVs, furniture, and other heavy items are secure so your child can't pull them over.
- Keep your child within arm's reach when he is near or in water.
- Empty buckets, pools, and tubs when you are finished using them.
- Never leave young brothers or sisters in charge of your child.
- When you go out, put a hat on your child, have him wear sun protection clothing, and apply sunscreen with SPF of 15 or higher on his exposed skin. Limit time outside when the sun is strongest (11:00 am–3:00 pm).
- Keep your child away when your pet is eating. Be close by when he plays with your pet.
- Keep poisons, medicines, and cleaning supplies in locked cabinets and out of your child's sight and reach.
- Keep cords, latex balloons, plastic bags, and small objects, such as marbles and batteries, away from your child. Cover all electrical outlets.
- Put the Poison Help number into all phones, including cell phones. Call if you are worried your child has swallowed something harmful. Do not make your child vomit.

WHAT TO EXPECT AT YOUR CHILD'S 15 MONTH VISIT

We will talk about

- Supporting your child's speech and independence and making time for yourself
- Developing good bedtime routines
- Handling tantrums and discipline
- Caring for your child's teeth
- Keeping your child safe at home and in the car

Consistent with *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*, 4th Edition

For more information, go to <https://brightfutures.aap.org>.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



The information contained in this handout should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances. Original handout included as part of the *Bright Futures Tool and Resource Kit*, 2nd Edition.

Inclusion in this handout does not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of the resources mentioned in this handout. Web site addresses are as current as possible but may change at any time.

The American Academy of Pediatrics (AAP) does not review or endorse any modifications made to this handout and in no event shall the AAP be liable for any such changes.

© 2019 American Academy of Pediatrics. All rights reserved.

Choking Prevention and First Aid for Infants and Children



When children begin crawling, or eating table foods, parents must be aware of the dangers and risks of choking. Children younger than 5 years can easily choke on food and small objects.

Choking occurs when food or small objects get caught in the throat and block the airway. This can prevent oxygen from getting to the lungs and the brain. When the brain goes without oxygen for more than 4 minutes, brain damage or even death may occur. Many children die from choking each year. Most children who choke to death are younger than 5 years. Two thirds of choking victims are infants younger than 1 year.

Balloons, balls, marbles, pieces of toys, and foods cause the most choking deaths.

Read more about choking prevention and first aid.

Dangerous foods

Do not feed children younger than 4 years round, firm food unless it is chopped completely. Round, firm foods are common choking dangers. When infants and young children do not grind or chew their food well, they may try to swallow it whole. The following foods can be choking hazards:

- Hot dogs
- Nuts and seeds
- Chunks of meat or cheese
- Whole grapes
- Hard, gooey, or sticky candy
- Popcorn
- Chunks of peanut butter
- Raw vegetables
- Fruit chunks, such as apple chunks
- Chewing gum

Dangerous household items

Keep the following household items away from infants and children:

- Balloons
- Coins
- Marbles
- Toys with small parts
- Toys that can be squeezed to fit entirely into a child's mouth
- Small balls
- Pen or marker caps
- Small button-type batteries
- Medicine syringes

What you can do to prevent choking

- *Learn CPR (cardiopulmonary resuscitation) (basic life support).*
- *Be aware that balloons pose a choking risk to children up to 8 years of age.*
- *Keep the above foods from children until 4 years of age.*
- *Insist that children eat at the table, or at least while sitting down. They should never run, walk, play, or lie down with food in their mouths.*
- *Cut food for infants and young children into pieces no larger than one-half inch, and teach them to chew their food well.*
- *Supervise mealtime for infants and young children.*
- *Be aware of older children's actions. Many choking incidents occur when older brothers or sisters give dangerous foods, toys, or small objects to a younger child.*
- *Avoid toys with small parts, and keep other small household items out of the reach of infants and young children.*
- *Follow the age recommendations on toy packages. Age guidelines reflect the safety of a toy based on any possible choking hazard as well as the child's physical and mental abilities at various ages.*
- *Check under furniture and between cushions for small items that children could find and put in their mouths.*
- *Do not let infants and young children play with coins.*

First aid for the child who is choking

Make a point to learn the instructions on the reverse side of this brochure. Post the chart in your home. However, these instructions should *not* take the place of an approved class in basic first aid, CPR, or emergency prevention. Contact your local American Red Cross office or the American Heart Association to find out about classes offered in your area. Most of the classes teach basic first aid, CPR, and emergency prevention along with what to do for a choking infant or child. Your pediatrician also can help you understand these steps and talk to you about the importance of supervising mealtime and identifying dangerous foods and objects.

The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

From your doctor
IRL DOWELL PEDIATRICS

171 Franklin Tpke., Suite 110
Waldwick, NJ 07463

Phone: 201-612-5100 Fax: 201-612-4499

American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

The American Academy of Pediatrics is an organization of 60,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety, and well-being of infants, children, adolescents, and young adults.

American Academy of Pediatrics
Web site—www.aap.org

Copyright ©2006
American Academy of Pediatrics, Updated 4/06

