

# BRIGHT FUTURES HANDOUT ► PARENT

## 15 MONTH VISIT



Here are some suggestions from Bright Futures experts that may be of value to your family.

### TALKING AND FEELING

- Try to give choices. Allow your child to choose between 2 good options, such as a banana or an apple, or 2 favorite books.
- Know that it is normal for your child to be anxious around new people. Be sure to comfort your child.
- Take time for yourself and your partner.
- Get support from other parents.
- Show your child how to use words.
  - Use simple, clear phrases to talk to your child.
  - Use simple words to talk about a book's pictures when reading.
  - Use words to describe your child's feelings.
  - Describe your child's gestures with words.

### A GOOD NIGHT'S SLEEP

- Put your child to bed at the same time every night. Early is better.
- Make the hour before bedtime loving and calm.
- Have a simple bedtime routine that includes a book.
- Try to tuck in your child when he is drowsy but still awake.
- Don't give your child a bottle in bed.
- Don't put a TV, computer, tablet, or smartphone in your child's bedroom.
- Avoid giving your child enjoyable attention if he wakes during the night. Use words to reassure and give a blanket or toy to hold for comfort.

### TANTRUMS AND DISCIPLINE

- Use distraction to stop tantrums when you can.
- Praise your child when she does what you ask her to do and for what she can accomplish.
- Set limits and use discipline to teach and protect your child, not to punish her.
- Limit the need to say "No!" by making your home and yard safe for play.
- Teach your child not to hit, bite, or hurt other people.
- Be a role model.

### HEALTHY TEETH

- Take your child for a first dental visit if you have not done so.
- Brush your child's teeth twice each day with a small smear of fluoridated toothpaste, no more than a grain of rice.
- Wean your child from the bottle.
- Brush your own teeth. Avoid sharing cups and spoons with your child. Don't clean her pacifier in your mouth.

**Helpful Resources:** Poison Help Line: 800-222-1222

Information About Car Safety Seats: [www.safercar.gov/parents](http://www.safercar.gov/parents) | Toll-free Auto Safety Hotline: 888-327-4236

# 15 MONTH VISIT—PARENT

## SAFETY

- Make sure your child's car safety seat is rear facing until he reaches the highest weight or height allowed by the car safety seat's manufacturer. In most cases, this will be well past the second birthday.
- Never put your child in the front seat of a vehicle that has a passenger airbag. The back seat is the safest.
- Everyone should wear a seat belt in the car.
- Keep poisons, medicines, and lawn and cleaning supplies in locked cabinets, out of your child's sight and reach.
- Put the Poison Help number into all phones, including cell phones. Call if you are worried your child has swallowed something harmful. Don't make your child vomit.
- Place gates at the top and bottom of stairs. Install operable window guards on windows at the second story and higher. Keep furniture away from windows.
- Turn pan handles toward the back of the stove.
- Don't leave hot liquids on tables with tablecloths that your child might pull down.
- Have working smoke and carbon monoxide alarms on every floor. Test them every month and change the batteries every year. Make a family escape plan in case of fire in your home.

## WHAT TO EXPECT AT YOUR CHILD'S 18 MONTH VISIT

### We will talk about

- Handling stranger anxiety, setting limits, and knowing when to start toilet training
- Supporting your child's speech and ability to communicate
- Talking, reading, and using tablets or smartphones with your child
- Eating healthy
- Keeping your child safe at home, outside, and in the car

Consistent with *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents*, 4th Edition

For more information, go to <https://brightfutures.aap.org>.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®




The information contained in this handout should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances. Original handout included as part of the *Bright Futures Tool and Resource Kit*, 2nd Edition.

Inclusion in this handout does not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of the resources mentioned in this handout. Web site addresses are as current as possible but may change at any time.

The American Academy of Pediatrics (AAP) does not review or endorse any modifications made to this handout and in no event shall the AAP be liable for any such changes.

© 2019 American Academy of Pediatrics. All rights reserved.

# 1 to 2 Years



## 1 TO 2 YEARS

### Safety for Your Child

Did you know that injuries are the leading cause of death of children in the United States? Most of these injuries can be prevented.

Often, injuries happen because parents are not aware of what their children can do. At this age your child can *walk, run, climb, jump, and explore* everything. Because of all the new things he or she can do, this stage is a very dangerous time in your child's life. It is your responsibility to protect your child from injury. Your child cannot understand danger or remember "no" while exploring.

#### Firearm Hazards

Children in homes where guns are present are in more danger of being shot by themselves, their friends, or family members than of being injured by an intruder. It is best to keep all guns out of the home. **Handguns are especially dangerous.** If you keep a gun, keep it unloaded and in a locked place, with the ammunition locked separately. Ask if the homes where your child visits or is cared for have guns and how they are stored.

#### Poisonings

Children continue to explore their world by putting everything in their mouths, even if it doesn't taste good. Your child can *open doors and drawers, take things apart, and open bottles* easily now, so you must use safety caps on all medicines and toxic household products. **Keep the safety caps on** at all times or find safer substitutes to use. Contact Poison Help for more information.


Your child is now able to get into and on top of everything. Be sure to keep all household products and medicines completely out of sight and reach. Never store lye drain cleaners in your home. Keep all products in their original containers. Use medicines exactly as directed and dispose of unused medicine safely as soon as you are finished with it.

**If your child does put something poisonous into his or her mouth, call Poison Help immediately. Add the Poison Help line (1-800-222-1222) to your phone contacts list. Do not make your child vomit.**

#### Falls

To prevent serious falls, lock the doors to any dangerous areas. **Use gates on stairways and install operable window guards** above the first floor. **Remove sharp-edged furniture** from the room your child plays and sleeps in. At this age your child will walk well and start to climb, jump, and run as well. A chair left next to a kitchen counter, table, or window allows your child to climb to dangerously high places. Remember, your child does not understand what is dangerous.

**If your child has a serious fall or does not act normally after a fall, call your doctor.**



American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



(over)

## Burns

The kitchen is a dangerous place for your child during meal preparation. Hot liquids, grease, and hot foods spilled on your child will cause serious burns. A **safer place for your child** while you are cooking, eating, or unable to give him or her your full attention is the **playpen, crib, or stationary activity center, or buckled into a high chair**. It's best to keep your child out of the kitchen while cooking.



Children who are learning to walk will grab anything to steady themselves, including hot oven doors, wall heaters, or outdoor grills. Keep your child out of rooms where there are hot objects that may be touched, or put a barrier around them. If you have a gas fireplace, keep children away while it is in use and for at least an hour after turning it off. The glass doors get extremely hot and can cause severe burns.

Your child will *reach* for your hot food or cup of coffee, so don't leave it within your child's reach.

**NEVER carry your child and hot liquids at the same time.** You can't handle both.

**If your child does get burned, immediately put cold water on the burned area. Keep the burned area in cold water for a few minutes to cool it off. Then cover the burn loosely with a dry bandage or clean cloth. Call your doctor for all burns. To protect your child from tap water scalds, the hottest temperature at the faucet should be no more than 120°F. In many cases you can adjust your water heater.**

**Make sure you have a working smoke alarm on every level of your home, especially in furnace and sleeping areas. Test the alarms every month. It is best to use smoke alarms that use long-life batteries, but if you do not, change the batteries at least once a year.**

## Drowning

At this age your child loves to play in water. **NEVER leave your child alone in or near a bathtub, pail of water, wading or swimming pool, or any other water, even for a moment.** Empty all buckets after each use. Keep the bathroom doors closed. Your child can drown in less than 2 inches of water. Knowing how to swim does NOT mean your child is safe near or in water. Stay within an arm's length of your child around water.



If you have a swimming pool, fence it on all 4 sides with a fence at least 4 feet high, and be sure the gates are self-latching. If possible, lock doors that could lead to the pool area. Most children drown when they wander out of the house and fall into a pool that is not fenced off from the house. You cannot watch your child every minute while he or she is in the house. It only takes a moment for your child to get out of your house and fall into your pool.

## And Remember Car Safety

**Car crashes are a great danger to your child's life and health.** The crushing forces to your child's brain and body in a crash or sudden stop, even at low speeds, can cause severe injuries or death. **To prevent these injuries USE a car safety seat EVERY TIME your child rides in the car.** All infants and toddlers should ride in a rear-facing car safety seat until they reach the highest weight or height allowed by their car safety seat's manufacturer. Be sure that the safety seat is installed and used correctly. Read and follow the instructions that come with the car safety seat and the instructions for using car safety seats in the owners' manual of your car. **The safest place for all infants and children to ride is in the back seat.**



**Do not leave your child alone in or around the car. Keep vehicles and their trunks locked.** Children who are left in a car can die of heatstroke because temperatures can reach deadly levels in minutes. They can be strangled by power windows or knock the vehicle into gear.

Always walk behind your car to be sure your child is not there before you back out of your driveway. You may not see your child behind your car in the rearview mirror.

**Remember, the biggest threat to your child's life and health is an injury.**

From Your Doctor

**LIVE WELL PEDIATRICS**

171 Franklin Tpke., Suite 110


Waldwick, NJ 07463

Phone: 201-612-5100 Fax: 201-612-4499

The information in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

# Car Seat Recommendations for Children

There are many car seat choices on the market. Use the information below to help you choose the type of car seat that best meets your child's needs.



- Select a car seat based on your child's age and size, choose a seat that fits in your vehicle, and use it every time.
- Always refer to your specific car seat manufacturer's instructions (check height and weight limits) and read the vehicle owner's manual on how to install the car seat using the seat belt or lower anchors and a tether, if available.
- To maximize safety, keep your child in the car seat for as long as possible, as long as the child fits within the manufacturer's height and weight requirements.
- Keep your child in the back seat at least through age 12.

## Rear-Facing Car Seat

### Birth – 12 Months



Your child under age 1 should always ride in a rear-facing car seat. There are different types of rear-facing car seats:

- Infant-only seats can only be used rear-facing.
- Convertible and All-in-One car seats typically have higher height and weight limits for the rear-facing position, allowing you to keep your child rear-facing for a longer period of time.

### 1 – 3 Years



Keep your child rear-facing as long as possible. It's the best way to keep him or her safe. Your child should remain in a rear-facing car seat until he or she reaches the top height or weight limit allowed by your car seat's manufacturer. Once your child outgrows the rear-facing car seat, your child is ready to travel in a forward-facing car seat with a harness and tether.

## Forward-Facing Car Seat

### 1 - 3 Years

Keep your child rear-facing as long as possible. It's the best way to keep him or her safe. Your child should remain in a rear-facing car seat until he or she reaches the top height or weight limit allowed by your car seat's manufacturer. Once your child outgrows the rear-facing car seat, your child is ready to travel in a forward-facing car seat with a harness and tether.



### 4 - 7 Years

Keep your child in a forward-facing car seat with a harness and tether until he or she reaches the top height or weight limit allowed by your car seat's manufacturer. Once your child outgrows the forward-facing car seat with a harness, it's time to travel in a booster seat, but still in the back seat.



---

## Booster Seat

### 4 - 7 Years

Keep your child in a forward-facing car seat with a harness and tether until he or she reaches the top height or weight limit allowed by your car seat's manufacturer. Once your child outgrows the forward-facing car seat with a harness, it's time to travel in a booster seat, but still in the back seat.



### 8 - 12 Years

Keep your child in a booster seat until he or she is big enough to fit in a seat belt properly. For a seat belt to fit properly the lap belt must lie snugly across the upper thighs, not the stomach. The shoulder belt should lie snug across the shoulder and chest and not cross the neck or face. Remember: your child should still ride in the back seat because it's safer there.



---

## Seat Belt

### 8 - 12 Years

Keep your child in a booster seat until he or she is big enough to fit in a seat belt properly. For a seat belt to fit properly the lap belt must lie snugly across the upper thighs, not the stomach. The shoulder belt should lie snug across the shoulder and chest and not cross the neck or face. Remember: your child should still ride in the back seat because it's safer there.



---

[NHTSA.gov/TheRightSeat](http://NHTSA.gov/TheRightSeat)



U.S. Department of Transportation  
National Highway Traffic Safety Administration

