

Claritin Chewable Tablet 24 Hour (Loratadine 5mg)

Age	Dose
children under 2 years of age	ask a doctor
children 2 years to under 6 years of age	chew 1 tablet daily; not more than 1 tablet in 24 hours
adults and children 6 years and over	chew 2 tablets daily; not more than 2 tablets in 24 hours
consumers with liver disease or kidney disease	ask a doctor

Children's Claritin Syrup 24-Hour (loratadine 5 mg in 5 mL)

Age	Dose
children under 2 years of age	ask a doctor
children 2 years to under 6 years of age	5 mL once daily; do not take more than 5 mL in 24 hours
adults and children 6 years and over	10 mL once daily; do not take more than 10 mL in 24 hours
consumers with liver disease or kidney disease	ask a doctor

Claritin RediTabs for Juniors 12-Hour (loratadine 5 mg)

Age	Dose
children under 6 years of age	ask a doctor
adults and children 6 years and over	1 tablet every 12 hours; not more than 2 tablets in 24 hours
consumers with liver disease or kidney disease	ask a doctor

Claritin RediTabs for Juniors 24-Hour (loratadine 10 mg)

Age	Dose
children under 6 years of age	ask a doctor
adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours
consumers with liver disease or kidney disease	ask a doctor